Mississippi Behavioral Risk Factor Surveillance System Trend Report, 1993-2002

<u>Note</u>: Unless otherwise indicated the denominator is the total sample population except that all respondents answering "unknown" or who refused to answer a question are excluded from the denominator.

| Indicator | Rates | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 |
|---|--------------|------|------|------|------|------|------|------|------|------|------|
| Health status | | | | | | | | | | | |
| Donort that any and health is fair any as | Actual | 21.6 | 22.1 | 21.4 | 21.1 | 21.9 | 21.0 | 20.9 | 20.2 | 23.0 | 25.2 |
| Report that general health is fair or poor | Age Adjusted | 20.9 | 20.5 | 21.8 | 21.2 | 22.1 | 21.1 | 20.9 | 20.3 | 23.1 | 24.8 |
| Report that physical health was not good for more than 7 days during past 30 days | Actual | 11.7 | 9.9 | 11.2 | 11.6 | 11.4 | 14.1 | 13.1 | 12.6 | 15.8 | |
| | Age Adjusted | 12.0 | 10.4 | 11.5 | 11.9 | 11.6 | 14.3 | 13.4 | 13.0 | 16.1 | |
| Report that mental health was not | Actual | 11.1 | 10.2 | 9.6 | 9.4 | 9.4 | 12.1 | 13.2 | 14.1 | 13.7 | |
| good for more than 7 days during past 30 days | Age Adjusted | 11.2 | 10.2 | 9.6 | 9.6 | 9.6 | 11.7 | 13.2 | 13.3 | 13.8 | |
| Report poor physical/mental health prevented | Actual | 7.7 | 4.7 | 7.8 | 6.9 | 5.5 | 7.4 | 8.3 | 7.9 | 9.5 | |
| usual activities for more than 7 days in past 30 days (Denominator: Total Sample) | Age Adjusted | 7.7 | 4.9 | 7.9 | 6.9 | 5.5 | 7.5 | 8.1 | 8.0 | 9.6 | |
| Report poor physical/mental health prevented usual activities for more than 7 days in past 30 | Actual | 15.7 | 11.1 | 19.8 | 18.5 | 13.8 | 16.7 | 17.3 | 17.0 | 20.3 | |
| days (Denominator: Excludes those with zero days of poor physical/mental health) | Age Adjusted | 16.8 | 12.0 | 20.4 | 18.9 | 14.3 | 14.2 | 17.9 | 18.1 | 20.9 | |
| Cigarette smoking | | | | | | | | | | | |
| Demont there are exament alcomette employee | Actual | 24.2 | 22.2 | 24.1 | 23.2 | 23.1 | 24.2 | 23.4 | 23.5 | 25.4 | 27.4 |
| Report they are current cigarette smokers | Age Adjusted | 24.5 | 22.6 | 24.4 | 23.3 | 23.5 | 24.5 | 23.4 | 23.6 | 25.4 | 27.2 |
| III ah sahaal advastion anlass | Actual | 28.6 | 26.8 | 34.5 | 29.8 | 29.5 | 30.1 | 30.3 | 32.2 | 31.2 | 36.3 |
| High school education or less | Age Adjusted | 34.6 | 36.5 | 41.5 | 37.3 | 36.7 | 36.2 | 37.1 | 37.1 | 35.8 | 41.2 |
| | Actual | 20.3 | 18.7 | 22.4 | 21.0 | 17.7 | 21.2 | 18.3 | 18.3 | 20.3 | 24.7 |
| Blacks | Age Adjusted | 21.8 | 20.1 | 23.9 | 20.7 | 19.3 | 22.1 | 17.9 | 18.8 | 19.8 | 24.3 |

| Indicator | Rates | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 |
|---|--------------|------|------|------|------|------|------|------|------|------|------|
| | Actual | 20.8 | 21.4 | 20.6 | 21.1 | 21.3 | 21.9 | 20.6 | 20.5 | 20.9 | 20.0 |
| Report that they are former cigarette smokers | Age Adjusted | 21.8 | 22.0 | 20.9 | 21.6 | 21.7 | 22.1 | 20.0 | 20.8 | 21.2 | 20.2 |
| Daily smokers who report that they have quit | Actual | 63.7 | 46.2 | 43.9 | 49.8 | 51.2 | 51.0 | 54.6 | 46.7 | 51.5 | 53.9 |
| smoking for 1 day or more in the past 12 months | Age Adjusted | 64.1 | 40.7 | 43.2 | 47.6 | 50.4 | 52.1 | 54.7 | 47.0 | 49.9 | 52.0 |
| Quit ratio (Ever smokers who are former | Actual | 46.2 | 49.0 | 46.1 | 47.6 | 47.9 | 47.5 | 46.9 | 46.5 | 45.1 | 42.2 |
| smokers) | Age Adjusted | 44.9 | 48.0 | 45.2 | 46.8 | 46.1 | 46.3 | 44.5 | 46.1 | 44.3 | 42.0 |
| Alcohol use during the past month | | | | | | | | | | | |
| Panart aguta (hinga) drinking | Actual | 8.7 | | 8.8 | | 9.6 | | 10.9 | | 11.7 | 12.4 |
| Report acute (binge) drinking | Age Adjusted | 8.3 | | 8.5 | | 9.4 | | 11.5 | | 11.5 | 12.3 |
| Depart charge deinling | Actual | 2.8 | | 2.7 | | 2.5 | | 3.3 | | 4.5 | 4.6 |
| Report chronic drinking | Age Adjusted | 2.6 | | 2.8 | | 2.4 | | 3.5 | | 4.4 | 4.6 |
| Physical activity/exercise | | | | | | | | | | | |
| Report that they are physically inactive | Actual | | 38.5 | | 39.6 | | 33.8 | | 33.4 | | |
| Report that they are physically mactive | Age Adjusted | | 39.1 | | 39.7 | | 34.1 | | 33.6 | | |
| 65+ year olds (No age adjustment) | Actual | | 51.1 | | 49.2 | | 40.2 | | 43.1 | | |
| Income less than \$20,000 | Actual | | 47.0 | | 49.0 | | 44.0 | | 42.3 | | |
| income less than \$20,000 | Age Adjusted | | 47.6 | | 48.0 | | 45.0 | | 43.7 | | |
| Report sedentary lifestyle (sedentary or | Actual | | 67.1 | | 64.7 | | 61.2 | | 61.1 | | |
| irregular physical activity profile) | Age Adjusted | | 67.6 | | 64.6 | | 61.4 | | 61.4 | | |
| Deposit frequent recorder consists | Actual | | 14.0 | | 17.0 | | 19.1 | | 18.6 | | |
| Report frequent regular exercise | Age Adjusted | | 13.7 | | 16.9 | | 18.9 | | 18.5 | | |

| Indicator | Rates | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 |
|---|--------------|------|------|------|------|------|------|------|------|------|------|
| | Actual | | 9.8 | | 11.2 | | 10.4 | | 9.4 | | |
| Report regular intensive exercise | Age Adjusted | | 9.9 | | 11.5 | | 10.4 | | 9.6 | | |
| In a great land them \$20,000 | Actual | | 6.7 | | 8.4 | | 8.4 | | 6.6 | | |
| Income less than \$20,000 | Age Adjusted | | 6.4 | | 8.0 | | 7.6 | | 6.0 | | |
| Report using physical activity/exercise to lose weight or maintain weight | Actual | | 34.4 | | 31.2 | | 39.5 | | 40.7 | | |
| Denominator: Total Sample | Age Adjusted | | 34.3 | | 31.3 | | 39.5 | | 40.6 | | |
| Report using physical activity/exercise to lose or | Actual | | 57.8 | | 42.0 | | 52.4 | | 54.6 | | |
| maintain weight. (Denominator: Those trying to lose or maintain weight) | Age Adjusted | | 56.0 | | 41.7 | | 52.3 | | 54.2 | | |
| Body weight and weight control | | | | | | | | | | | |
| Decreased acts who are suggested (DML > 25.0) | Actual | 54.6 | 53.8 | 54.7 | 54.5 | 57.9 | 59.1 | 62.3 | 61.8 | 61.1 | 63.1 |
| Respondents who are overweight (BMI ≥25.0) | Age Adjusted | 55.2 | 54.6 | 55.3 | 55.4 | 58.2 | 59.8 | 62.0 | 62.1 | 61.5 | 63.6 |
| Respondents who are obese (BMI ≥30) | Actual | 18.9 | 19.7 | 19.6 | 19.8 | 22.0 | 22.9 | 23.4 | 25.0 | 25.4 | 26.8 |
| Respondents who are obese (BMT 250) | Age Adjusted | 19.4 | 20.5 | 19.8 | 20.2 | 22.2 | 23.1 | 23.2 | 25.3 | 25.7 | 27.0 |
| Report they are dieting to lose or maintain weight | Actual | | 54.9 | | 52.4 | | 54.5 | | 55.6 | | |
| (Denominator : Total Sample) | Age Adjusted | | 55.7 | | 52.8 | | 55.0 | | 55.8 | | |
| Report they are dieting to lose or maintain weight | Actual | | 72.3 | | 71.3 | | 71.5 | | 74.1 | | |
| (Denominator : Those trying to lose or maintain weight) | Age Adjusted | | 72.3 | | 70.2 | | 71.7 | | 74.3 | | |
| Report using physical activity/exercise to lose or | Actual | | 34.4 | | 31.2 | | 39.5 | | 40.7 | | |
| maintain weight. (Denominator: Total sample) | Age Adjusted | | 34.4 | | 31.3 | | 39.5 | | 40.6 | | |
| Report using physical activity/exercise to lose or | Actual | | 45.2 | | 42.0 | | 52.4 | | 54.6 | | |
| maintain weight. (Denominator: Those trying to lose or maintain weight) | Age Adjusted | | 45.0 | | 41.7 | | 52.3 | | 54.2 | | |

| Indicator | Rates | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 |
|--|--------------|------|------|------|------|------|------|------|------|------|------|
| At risk of developing diabetes | | | | | | | | | | | |
| Those at risk of developing type 2 diabetes | Actual | | 22.6 | | 24.1 | | 24.4 | | 26.1 | 24.3 | 24.5 |
| because of weight or lack of exercise | Age Adjusted | | 24.5 | | 25.5 | | 25.3 | | 26.8 | 24.7 | 24.4 |
| Diet/Nutrition | | | | | | | | | | | |
| Report that they eat fruits/vegetables | Actual | | 13.9 | | 18.2 | | 15.6 | | 18.5 | | 19.1 |
| 5 or more times per day | Age Adjusted | | 14.2 | | 18.1 | | 15.6 | | 18.5 | | 19.1 |
| Cholesterol awareness | | | | | | | | | | | |
| Report that they have ever | Actual | 62.3 | | 62.0 | | 66.5 | | 72.9 | | 73.8 | |
| had a blood cholesterol test | Age Adjusted | 63.1 | | 62.9 | | 67.2 | | 71.4 | | 74.2 | |
| Those tested who have had blood | Actual | 45.5 | | 43.6 | | 46.0 | | 49.0 | | 51.8 | |
| cholesterol checked within the past year | Age Adjusted | 46.4 | | 44.3 | | 46.5 | | 47.9 | | 52.0 | |
| Report blood cholesterol checked | Actual | 55.1 | | 53.5 | | 55.1 | | 60.9 | | 62.4 | |
| within the past 2 years | Age Adjusted | 55.9 | | 54.3 | | 55.6 | | 59.7 | | 62.7 | |
| Report blood cholesterol | Actual | 60.2 | | 58.3 | | 62.5 | | 67.0 | | 69.2 | |
| checked within the past 5 years | Age Adjusted | 61.1 | | 59.1 | | 63.2 | | 65.7 | | 68.6 | |
| Report being told by a health professional | Actual | 19.8 | | 15.8 | | 19.2 | | 22.5 | | 22.8 | |
| that they have high blood cholesterol | Age Adjusted | 20.5 | | 16.2 | | 19.6 | | 22.4 | | 23.0 | |
| Diabetes | | | | | | | | | | | |
| Report being told by a doctor | Actual | 6.4 | 5.7 | 6.3 | 6.1 | 6.1 | 7.7 | 8.3 | 7.6 | 9.3 | 8.6 |
| that they have diabetes | Age Adjusted | 6.5 | 5.9 | 6.4 | 6.2 | 6.3 | 7.7 | 8.0 | 7.7 | 9.3 | 8.7 |
| Those with diabetes who have had two or more visits to a health professional for diabetes in the | Actual | | | | 74.4 | 74.7 | 75.7 | | 80.7 | | 80.4 |
| past year | Age Adjusted | | | | 59.9 | 72.2 | 68.6 | | 79.6 | | 81.0 |

| Indicator | Rates | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | |
|--|--------------|------|------|------|------|------|------|------|------|------|------|--|
| Those with diabetes who have had at least four | Actual | | | | 59.1 | 56.5 | 46.8 | | 51.0 | | 51.0 | |
| visits to a health professional for their diabetes in past year | Age Adjusted | | | | 48.5 | 51.0 | 43.1 | | 49.8 | | 42.9 | |
| Those with diabetes who have had two or more | Actual | | | | 29.3 | 43.5 | 38.4 | | 38.6 | | 55.4 | |
| foot exams by a health professional in past year | Age Adjusted | | | | 23.3 | 49.3 | 32.4 | | 38.4 | | 46.4 | |
| Those with diabetes who have had at least four | Actual | | | | 23.3 | 29.9 | 24.8 | | 19.7 | | 28.0 | |
| foot exams by a health professional in past year | Age Adjusted | | | | 18.3 | 24.5 | 19.6 | | 21.4 | | 25.4 | |
| Those with diabetes who have had | Actual | | | | 55.4 | 50.2 | 54.8 | | 56.5 | | 66.5 | |
| a dilated eye exam within the past year | Age Adjusted | | | | 45.6 | 45.3 | 53.3 | | 47.6 | | 62.8 | |
| Those with diabetes who have had | Actual | | | | 69.6 | 70.3 | 68.1 | | 66.3 | | 80.3 | |
| a dilated eye exam within the past 2 years | Age Adjusted | | | | 54.0 | 66.5 | 63.3 | | 55.7 | | 80.9 | |
| Mammography screening (Age Adjustment Not Applicable) | | | | | | | | | | | | |
| Report that they have ever had a CBE and mammogram (Women 40+) | Actual | 64.1 | 64.7 | 67.2 | 71.9 | 71.8 | 73.1 | 72.6 | 77.7 | 78.1 | 77.9 | |
| Income less than \$10,000 | Actual | 52.4 | 50.0 | 51.6 | 55.5 | 68.6 | 74.9 | 56.2 | 62.8 | 69.9 | 60.6 | |
| Less than a high school education | Actual | 54.2 | 49.3 | 50.2 | 57.3 | 59.5 | 55.9 | 61.2 | 61.6 | 66.4 | 66.7 | |
| Women aged 70+ years of age | Actual | 59.7 | 53.0 | 57.5 | 61.8 | 62.0 | 63.7 | 69.3 | 80.9 | 72.6 | 74.4 | |
| Black women | Actual | 59.4 | 59.4 | 49.8 | 62.3 | 61.1 | 67.1 | 63.7 | 67.8 | 71.2 | 69.4 | |
| Report having a CBE and mammogram in past 2 years (Women 50+) | Actual | 51.4 | 49.5 | 49.1 | 53.8 | 60.7 | 58.4 | 56.6 | 61.6 | 60.9 | 62.3 | |
| Income less than \$10,000 | Actual | 41.0 | 32.6 | 37.8 | 38.3 | 60.4 | 45.6 | 40.0 | 45.7 | 52.6 | 44.1 | |
| Less than a high school education | Actual | 39.0 | 34.3 | 35.8 | 39.3 | 50.0 | 41.7 | 38.2 | 42.8 | 47.9 | 50.7 | |
| Women aged 70+ years of age | Actual | 47.9 | 38.9 | 45.5 | 49.0 | 50.7 | 47.7 | 49.8 | 57.7 | 53.5 | 53.7 | |
| Black women | Actual | 43.6 | 43.4 | 35.0 | 47.7 | 49.3 | 56.5 | 47.4 | 54.9 | 55.1 | 56.5 | |
| Report having a screening mammogram within the past year (Women 50-59) | Actual | 43.6 | 45.2 | 45.4 | 42.4 | 52.8 | 53.5 | 51.1 | 58.5 | 60.9 | 55.8 | |

| Indicator | Rates | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | |
|--|--------------|------|------|------|------|------|------|------|------|------|------|--|
| Report having a screening mammogram within the past 2 years (Women 50-59) | Actual | 60.3 | 62.3 | 56.3 | 64.2 | 68.6 | 71.9 | 65.6 | 73.1 | 75.2 | 72.8 | |
| Report having screening mammogram within the past year (Women 60-69) | Actual | 45.6 | 47.1 | 46.1 | 48.0 | 55.7 | 48.4 | 49.7 | 49.0 | 55.5 | 60.7 | |
| Report having a screening mammogram within the past 2 years (Women 60-69) | Actual | 60.0 | 58.5 | 56.7 | 65.0 | 69.8 | 69.3 | 68.6 | 67.1 | 69.8 | 73.7 | |
| Pap smear screening | | | | | | | | | | | | |
| Report that they have ever had a pap smear test | Actual | 93.1 | 94.1 | 92.6 | 92.7 | 95.8 | 93.3 | 94.6 | 95.5 | 96.5 | 90.1 | |
| (Women 18+ years of age who have never had a hysterectomy) | Age Adjusted | 92.5 | 93.3 | 93.1 | 92.5 | 95.8 | 93.7 | 95.1 | 95.2 | 96.7 | 90.7 | |
| Lucama lass than \$10,000 | Actual | 88.3 | 86.5 | 88.8 | 82.2 | 89.5 | 93.5 | 93.6 | 86.9 | 92.9 | 84.1 | |
| Income less than \$10,000 | Age Adjusted | 91.4 | 87.3 | 89.9 | 86.9 | 92.8 | 92.9 | 95.6 | 88.1 | 95.0 | 84.7 | |
| Lagrathan a high sahaal advastion | Actual | 84.7 | 83.5 | 85.5 | 88.3 | 90.9 | 85.7 | 87.8 | 86.8 | 94.9 | 86.7 | |
| Less than a high school education | Age Adjusted | 87.8 | 85.8 | 90.3 | 92.1 | 94.0 | 90.5 | 92.1 | 87.3 | 96.3 | 88.2 | |
| Women aged 70+ years of age (Age Adjustment not Applicable) | Actual | 81.1 | 79.6 | 80.9 | 79.3 | 82.1 | 86.8 | 85.0 | 85.6 | 91.9 | 82.1 | |
| Report that they have had a pap smear test | Actual | 79.6 | 84.4 | 83.4 | 81.8 | 85.8 | 83.4 | 83.0 | 87.0 | 86.3 | 82.7 | |
| in the past 3 years (Women 18+ years of age who have never had a hysterectomy) | Age Adjusted | 77.1 | 81.3 | 81.4 | 79.2 | 83.6 | 82.4 | 80.9 | 84.3 | 84.2 | 80.3 | |
| | Actual | 70.9 | 72.9 | 75.2 | 65.5 | 73.4 | 77.5 | 70.7 | 68.1 | 79.1 | 72.5 | |
| Income less than \$10,000 | Age Adjusted | 72.3 | 73.1 | 81.4 | 69.6 | 78.1 | 75.4 | 70.0 | 68.6 | 76.8 | 72.4 | |
| | Actual | 63.1 | 67.3 | 70.5 | 67.1 | 74.3 | 62.3 | 66.4 | 74.4 | 70.0 | 69.5 | |
| Less than a high school education | Age Adjusted | 69.8 | 69.5 | 77.7 | 73.5 | 71.3 | 69.3 | 74.9 | 74.5 | 72.2 | 74.0 | |
| Women aged 70+ years of age (Age Adjustment Not Applicable) | Actual | 51.1 | 58.7 | 51.7 | 50.2 | 55.1 | 63.9 | 52.6 | 75.5 | 64.4 | 46.1 | |
| Report that they have had pap smear test | Actual | 85.0 | 87.6 | 87.1 | 86.8 | 89.5 | 86.5 | 87.4 | 89.5 | 89.3 | 86.4 | |
| in the past 3 years (Women 18-64) | Age Adjusted | 81.8 | 84.8 | 85.8 | 84.7 | 87.9 | 86.4 | 86.2 | 87.2 | 88.5 | 85.2 | |

| Indicator | Rates | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 |
|--|------------------|--------|--------|---------|----------|------|------|------|------|------|------|
| Colorectal cancer screening: Persons aged 50+ y | ears of age (Age | Adjust | ment N | ot Appl | licable) | | | | | 84.7 | 81.8 |
| Report that they have ever had a sigmoidoscopy/proctoscopy exam | Actual | 33.6 | | 36.3 | | 35.8 | | 36.4 | | 40.8 | 42.9 |
| Report having a sigmoidoscopy or proctoscopy exam in the past year | Actual | 9.5 | | 12.5 | | 10.0 | | 12.5 | | 13.4 | 13.4 |
| Report having a sigmoidoscopy or proctoscopy exam in past 2 years | Actual | 14.3 | | 20.4 | | 17.7 | | 18.9 | | 21.6 | 21.9 |
| Report having had a sigmoidoscopy proctoscopy exam in past 5 years | Actual | 21.7 | | 27.3 | | 25.1 | | 27.0 | | 31.9 | 33.4 |
| Hypertension awareness | | | | | | | | | | | |
| Report being told by a health professional | Actual | 30.0 | | 29.8 | 29.5 | 34.3 | | 35.0 | | 31.4 | |
| that they have high blood pressure | Age Adjusted | 29.9 | | 30.2 | 29.9 | 34.6 | | 33.8 | | 31.7 | |
| Report having blood pressure taken by a | Actual | 88.3 | | 86.6 | 89.4 | 90.2 | | 89.4 | | | |
| health professionalin the past year | Age Adjusted | 88.4 | | 86.6 | 89.2 | 90.1 | | 89.3 | | | |
| Report having pressure taken by a | Actual | 93.5 | | 92.0 | 95.6 | 95.6 | | 96.0 | | | |
| health professional in the past 2 years | Age Adjusted | 93.5 | | 92.0 | 95.5 | 95.5 | | 96.0 | | | |
| Report having blood pressure taken by a | Actual | 97.2 | | 95.9 | 98.0 | 97.9 | | 98.3 | | | |
| health professional in the past 5 years | Age Adjusted | 97.1 | | 95.5 | 97.9 | 97.9 | | 98.3 | | | |

| Indicator | Rates | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 |
|--|--------------|------|------|------|------|------|------|------|------|------|------|
| Immunization | | | | | | | | | | | |
| Report having had a flu shot | Actual | 19.3 | | 26.3 | | 28.6 | | 31.3 | | 29.2 | 30.6 |
| within the past 12 months | Age Adjusted | 19.2 | | 26.1 | | 23.2 | | 30.6 | | 29.0 | 30.7 |
| Persons 65+ years of age (Age Adjustment Not Applicable) | Actual | 43.0 | | 57.0 | | 61.1 | | 62.9 | | 61.8 | 63.0 |
| | Actual | 14.0 | | 15.0 | | 18.2 | | 17.0 | | 21.2 | 19.9 |
| Report that having had a pneumonia vaccination | Age Adjusted | 13.4 | | 14.6 | | 17.6 | | 16.5 | | 20.6 | 19.8 |
| Persons aged 65+ years of age (Age Adjustment Not Applicable) | Actual | 27.8 | | 39.4 | | 45.9 | | 49.3 | | 55.7 | 58.9 |
| Health care access | | | | | | | | | | | |
| Report that they have | Actual | 16.1 | 12.7 | 14.0 | 14.2 | 15.1 | 18.8 | 16.9 | 19.3 | 18.7 | 22.8 |
| no kind of health care coverage | Age Adjusted | 15.7 | 13.7 | 14.0 | 14.3 | 15.1 | 18.6 | 17.4 | 19.0 | 18.6 | 22.5 |
| Report being unable to see doctor | Actual | 18.1 | 16.1 | 14.8 | 15.0 | 15.3 | 15.6 | 14.4 | 16.2 | | |
| in past 12 months because of cost | Age Adjusted | 18.2 | 16.4 | 14.9 | 14.6 | 15.3 | 15.6 | 14.5 | 16.2 | | |