

Status of Selected Year 2000 National Health Objectives for the State of Mississippi, 1997

Objective Number	Objective	Mississippi 1997 Status	Year 2000 Objective (National)
1.	Physical Activity and Fitness		
1.1	Coronary heart disease deaths	129.4	100.0 per 100,000 ¹
1.1a	Blacks	168.1	115.0 per 100,000 ¹
1.2	Reduce overweight (based on body mass index)	34	20.0 percent
1.3	Light to moderate physical activity (1996 data)	17	30.0 percent
1.4	Vigorous physical activity (1996 data)	11	20.0 percent
1.4a	Income less than \$20,000 (1996 data)	8	12.0 percent
1.5	No leisure-time activity (1996 data)	40	15.0 percent
1.5a	65+ year olds (1996 data)	49	22.0 percent
1.5c	Income less than \$20,000 (1996 data)	49	17.0 percent
2.	Nutrition		
2.2	Cancer deaths	140.7	130.0 per 100,000 ¹
3.	Tobacco		
3.2	Lung cancer deaths	45.7	42.0 per 100,000 ¹
3.3	Chronic obstructive pulmonary disease deaths	22.5	25.0 per 100,000 ¹
3.4	Cigarette smoking	23	15.0 percent
3.4a	High school education or less	28	20.0 percent
3.4d	Blacks	18	18.0 percent
4.	Alcohol and Other Drugs		
4.1	Alcohol-related motor vehicle crash deaths	30.7	8.5 per 100,000 ¹
4.1b	15-24 year olds	47.4	18.0 per 100,000
4.2	Cirrhosis deaths	7.6	6.0 per 100,000 ¹
4.2a	Black males	9.1	12.0 per 100,000 ¹
4.3	Drug-related deaths	2.3	3.0 per 100,000 ¹
5.	Family Planning		
5.1	Pregnancies 15-17 years old	61.3	50.0 per 1,000 females
5.1a	Blacks 15-19 years old	113.1	120.0 per 1,000 females
6.	Mental Health and Mental Disorders		
6.1	Suicides	11.6	10.5 per 100,000 ¹
6.1a	15-19 year olds	11.8	8.2 per 100,000
6.1b	Men 20-34 years old	22.5	21.4 per 100,000
6.1c	White men 65+ years old	55.9	39.2 per 100,000
7.	Violent and Abusive Behavior		
7.1	Homicides	14.6	7.2 per 100,000 ¹
7.1a	# 3 years old	6.4	3.1 per 100,000
7.1c	Black males 15-34 years old	82.2	72.4 per 100,000
7.1e	Black females 15-34 years old	14.7	16.0 per 100,000
7.3	Weapons-related violent deaths	22.2	12.6 per 100,000 ¹

Status of Selected Year 2000 National Health Objectives for the State of Mississippi, 1997

Objective Number	Objective	Mississippi 1997 Status	Year 2000 Objective (National)
9.	Unintentional Injuries		
9.1a	Unintentional injury deaths	50.6	29.3 per 100,000 ¹
9.1b	Black males	85.9	51.9 per 100,000
9.1c	White males	74.0	42.9 per 100,000
9.3	Motor vehicle crash deaths	31.0	16.8 per 100,000 ¹
9.3a	# 14 years	9.8	5.5 per 100,000
9.3b	15-24 years	47.6	33.0 per 100,000
9.3c	70+ years	46.2	20.0 per 100,000
9.3e	Motorcyclists	0.4	1.5 per 100,000 ¹
9.3f	Pedestrians	2.4	2.7 per 100,000 ¹
9.4	Falls and fall-related deaths	3.3	2.3 per 100,000 ¹
9.4a	65-84 years	26.6	14.4 per 100,000
9.4b	85+ years	177.8	105.0 per 100,000
9.4c	Black males 30-69 years old	7.0	5.6 per 100,000
9.5	Drowning deaths	2.7	1.3 per 100,000 ¹
9.5a	# 4 year olds	3.0	2.3 per 100,000
9.5b	Males 15-34 years old	5.9	2.5 per 100,000
9.5c	Black males	5.6	3.6 per 100,000
9.6	Fire deaths	3.3	1.2 per 100,000 ¹
9.6a	# 4 year olds	10.9	3.3 per 100,000
9.6b	65+ year olds	11.7	3.3 per 100,000
9.6c	Black males	7.6	4.3 per 100,000
9.6d	Black females	7.7	2.6 per 100,000
9.12	Safety belt use	57	85.0 percent
13.	Oral Health		
13.7	Oral cavity and pharynx cancer deaths		
	Males 45-74 years	12.1	10.5 per 100,000
	Females 45-74 years	3.3	4.1 per 100,000
14.	Maternal and Infant Health		
14.1	Infant mortality rate	10.6	7.0 per 1,000 LB ²
14.1a	Blacks	14.8	11.0 per 1,000 LB
14.1d	Neonatal mortality rate	6.5	4.5 per 1,000 LB
14.1e	Blacks	9.0	7.0 per 1,000 LB
14.1g	Postneonatal mortality rate	4.1	2.5 per 1,000 LB
14.1h	Blacks	5.8	4.0 per 1,000 LB
14.2	Fetal deaths	12.4	5.0 per 1,000 LB + FD ³
14.2a	Blacks	18.4	7.5 per 1,000 LB
14.3	Maternal mortality rate	26.5	3.3 per 100,000 LB
14.3a	Blacks	36.9	5.0 per 100,000 LB
14.5	Low birth weight (< 2,500 g)	10.1	5.0 percent of LB
14.5a	Blacks	13.4	9.0 percent of LB

Status of Selected Year 2000 National Health Objectives for the State of Mississippi, 1997

Objective Number	Objective	Mississippi 1997 Status	Year 2000 Objective (National)
	Very low birth weight (< 1,500 g)	2.0	1.0 percent of LB
	Blacks	2.8	2.0 percent of LB
14.8	Cesarean deliveries	26.1	15.0 percent of LB + FD ³
14.8a	Primary	17.4	12.0 percent of LB + FD ³
14.8b	Repeat	83.3	65.0 percent of LB + FD ^{3,4}
14.10	Abstinence from tobacco use	87.2	90.0 percent of LB
	Abstinence from alcohol use	99.0	95.0 percent of LB
14.11	Prenatal care-1st trimester	79.7	90.0 percent of LB
14.11a	Blacks	69.1	90.0 percent of LB
15.	Stroke		
15.2	Stroke deaths	35.1	20.0 per 100,000 ¹
15.2a	Blacks	52.8	27.0 per 100,000 ¹
15.14	Cholesterol checked within the past 5 years	60	75.0 percent
16.	Cancer		
16.3	Breast cancer deaths	20.0	20.6 per 100,000 females ¹
16.4	Uterine cervix cancer deaths	3.2	1.3 per 100,000 females ¹
16.5	Colorectal cancer deaths	12.3	13.2 per 100,000 ¹
16.11	Ever received a CBE ⁵ and mammogram (Women 40+)	71	80.0 percent
16.11b	Income less than \$10,000	68	80.0 percent
16.11c	Less than a high school education	59	80.0 percent
16.11d	Women aged 70+	61	80.0 percent
16.11e	Black women	61	80.0 percent
16.11	CBE ⁵ and mammogram in past 2 yrs (Women 50+)	60	60.0 percent
16.11b	Income less than \$10,000	60	60.0 percent
16.11c	Less than a high school education	50	60.0 percent
16.11d	Women aged 70+	49	60.0 percent
16.11e	Black women	48	60.0 percent
16.12	Ever received a pap test (Women aged 18+) ⁶	96	95.0 percent
16.12b	Women aged 70+	82	95.0 percent
16.12c	Less than a high school education	91	95.0 percent
16.12d	Income less than \$10,000	90	95.0 percent
16.12	Pap test in the past 3 yrs (Women aged 18+) ⁶	86	85.0 percent
16.12b	Women aged 70+	55	70.0 percent
16.12c	Less than a high school education	75	75.0 percent
16.12d	Income less than \$10,000	74	80.0 percent
16.13	Ever had a proctosigmoidoscopy (Persons aged 50+)	35	40.0 percent
17.	Diabetes and Chronic Disabling Conditions		
17.9	Diabetes-related deaths	50.2	38.0 per 100,000 ¹
	Blacks	82.7	58.0 per 100,000 ¹

Status of Selected Year 2000 National Health Objectives for the State of Mississippi, 1997

Objective Number	Objective	Mississippi 1997 Status	Year 2000 Objective (National)
19.	Sexually Transmitted Disease		
19.1	Gonorrhea	337.5	225.0 per 100,000
19.3	Primary and secondary syphilis	14.2	10.0 per 100,000
	Immunization and Infectious Disease		
20.3	Hepatitis A	3.2	23.0 per 100,000
20.4	Incidence of tuberculosis	9.0	3.5 per 100,000
20.7	Bacterial meningitis	2.6	4.7 per 100,000
20.11	Influenza immunization in the past 12 mos. (Persons aged 65+)	61	60.0 percent
	Ever had a pneumococcal immunization (Persons aged 65+)	44	60.0 percent
	Age-related Objectives		
	Death rate for children 1-14 years old	36.3	28.0 per 100,000 children 1-14
	Death rate for adolescents and young adults 15-24	131.2	85.0 per 100,000 people 15-24
	Death rate for adults 25-64	524.9	340.0 per 100,000 people 25-64

¹ Current and target rates are age-adjusted to the 1940 U.S. census population

² Live Births

³ Live Births + Fetal Deaths

⁴ Percentage of women who had previous Cesarean sections

⁵ Clinical breast exam

⁶ Women with an intact uterine cervix

Technical Note: Population-based rates were calculated using 1997 population estimates for Mississippi produced by the U.S. Bureau of the Census. Previous versions of this publication were based on the 1990 census population data.

For this reason, the population-based rates presented here are not directly comparable to those from previous years. Because the population estimates show Mississippi's population to be increasing, many of the rates will have decreased dramatically from those published last year. However, due to the changing age structure of the population, some rates (especially for younger age groups) may increase because of the use of 1997 population figures.

Sources: Healthy People 2000: National Health Promotion and Disease Prevention Objectives. PHS, U.S. Department of Health and Human Services, September 1990; Vital Statistics, 1997. Mississippi State Department of Health, Bureau of Public Health Statistics; "Mississippi Morbidity Report", Office of Epidemiology, Mississippi State Department of Health; and the Behavioral Risk Factor Surveillance System.