

2018-2019 Influenza Surveillance Report Week 43

Oct. 21 – Oct. 27, 2018


About our flu activity reporting

MSDH relies upon selected sentinel health practitioners across the state to report the percentage of total patient visits consistent with an influenza-like illness (ILI: fever of 100°F or higher AND cough and/or sore throat). Also, providers are supplied with specimen collection kits. Samples are submitted to the Mississippi Public Health Laboratory for influenza PCR testing. Reports are used to estimate the state's ILI rate and the magnitude of the state's influenza activity. Reports represent only the distribution of flu in the state, not an actual count of all flu cases statewide. *Information is provisional only and may change depending on additional reporting from sentinel providers*.

Content

- * <u>State ILI Surveillance</u>
- * Influenza Outbreaks
- * Flu Testing Reports
- * National and Mississippi Pediatric Mortality Surveillance
- * National ILI Surveillance
- * <u>Appendix</u>
 - **<u>Figure 1</u>** (Statewide ILI Data, Current Season compared to Previous Season)
 - Figure 2 (Percentage of ILI Cases by Age Group, Mississippi)
 - Figure 3 (Mississippi ILI Rates, 2018-2019 and Previous Seasons)
 - Figure 4 (State ILI Rates 2014-2018 (YTD))
 - Figure 5 (Comparison of the BioSense and Statewide ILI Rates)
 - Figure 6 (Comparison of Nationwide and Region 4 ILI Rates to Mississippi ILI Rates)
 - Figure 7 (CDC FluView Weekly Influenza Activity Map)

State ILI Surveillance




Total number of patients treated by sentinel providers in the last three weeks.| Table 1

2018-2019 Influenza Season					
CDC Week	Week Ending	Number of reports received from Sentinel Providers	Total patients	ILI symptoms	ILI Rate (%)
43	Oct. 27	158	16598	418	2.5
42	Oct. 20	157	17959	397	2.2
41	Oct. 13	156	17246	351	2.0


During week **43**, four districts (1, 4, 6, and 7) had an increase in ILI activity. The remaining five districts (2, 3, 5, 8, and 9) remained about the same. *Information is provisional only and may change depending on additional reporting from sentinel providers*. | **Table 2**




MSDH District ILI Rates (%) 2018-2019				
District	Week 42	Week 43		
State	2.2	2.5		
Ι	1.5	2.2		
II	2.4	1.9		
III	1.3	1.5		
IV	2.2	4.1		
V	2.3	1.9		
VI	1.9	3.4		
VII	2.7	4.0		
VIII	2.0	2.0		
IX	2.6	2.4		



The 2018-19 state ILI rate was **comparable** to the national and Region 4 baselines, but was **below** the state baseline, for week **43**. | Figure 3






Mississippi ILI Rates 2014-2018 | Figure 4

Syndromic ILI Surveillance

The Mississippi State Department of Health also collects influenza syndromic surveillance data through the CDC BioSense Platform. This data is comprised of chief complaints and diagnosis codes and is submitted electronically by participating hospitals and clinics throughout the state in near real-time. The BioSense data is an additional tool to monitor influenza activity in Mississippi.



has remained steady thus far in the 2018 – 2019 season, while the statewide ILI rate has slightly increased since week 41. | Figure 5

Influenza Outbreaks

Outbreaks are reportable in Mississippi as a Class 1A event and must be reported by telephone within **24 hours** of first knowledge or suspicion to the Mississippi State Department of Health. For more information on reportable diseases and conditions, please refer to the <u>MSDH List of Reportable Diseases and Conditions</u>.

No influenza outbreaks were reported to MSDH during week 43 (week ending October 27th).

For additional information on infection control measures in health care facilities and managing influenza outbreaks in long-term care facilities, please refer to the CDC's webpages: <u>https://www.cdc.gov/flu/professionals/infectioncontrol/index.htm</u> and <u>https://www.cdc.gov/flu/professionals/infectioncontrol/ltc-facility-guidance.htm</u>, respectively.

Flu Testing Reports

During week **43** (week ending October 27th), there were no laboratory confirmed influenza cases identified by MSDH Public Health Laboratory.

National and Mississippi Pediatric Mortality Surveillance

Nationally, **three** influenza-associated pediatric deaths were reported to CDC during week **43**. One death was associated with an influenza A(H1N1)pdm09 virus and occurred during week 41 (week ending October 13, 2018). **Two** influenza-associated pediatric deaths have been reported to CDC for the 2018-2019 season.


Two deaths that occurred during the 2017-2018 season were reported to CDC during week 43. One death was associated with an influenza A(H3) virus and one death was associated with an influenza B virus. These deaths bring the total number of reported influenza-associated deaths occurring during that season to **185**.

Mississippi has had no influenza-associated pediatric deaths reported during this influenza season.


For additional information on influenza-associated pediatric deaths, please refer to the CDC's FluView.

National ILI Surveillance

During week **43**, the Mississippi (2.5%), national (1.7%) and Region 4 (1.6%) ILI rates remained about the same as the previous week and were below their respective baselines. | Figure 6



During week **43**, influenza activity **remained low** in the United States, although small increases in activity were reported.¹ | <u>Figure 7</u>



¹For up-to-date information on flu activity nationwide, please refer to the CDC's website: <u>http://www.cdc.gov/flu/weekly/fluactivitysurv.htm</u>.

Level of Flu Activity	Definition
No Activity	Overall clinical activity remains low and there are no lab confirmed cases.
Sporadic	Isolated cases of lab confirmed influenza in the state; ILI activity is not increased <u>OR</u> A lab- confirmed outbreak in a single institution in the state; ILI activity is not increased.
Local	Increased ILI within a single region AND recent (within the past 3 weeks) laboratory evidence of influenza in that region. ILI activity in other regions is not increased <u>OR</u> two of more institutional outbreaks (ILI or lab confirmed) within a single region AND recent (within the past 3 weeks) lab confirmed influenza in that region. Other regions do not have increased ILI and virus activity is no greater than sporadic in those regions
Regional	Increased ILI in at least 2 regions but fewer than half of the regions AND recent (within the past 3 weeks) lab confirmed influenza in the affected regions <u>OR</u> Institutional outbreaks (ILI or lab confirmed) in at least 2 regions but fewer than half of the regions AND recent lab confirmed influenza in the affected regions.


Mississippi reported "No Activity" for the influenza activity during week 43. | Table 3

Widespread	Increased ILI and/or institutional outbreaks (ILI or lab confirmed) in at least half of the regions
	AND recent (within the past 3 weeks) lab confirmed influenza in the state.


Additional influenza information:

Centers for Disease Control and Prevention	http://cdc.gov/flu/
Centers for Disease Control and Prevention FluView	http://www.cdc.gov/flu/weekly/
MSDH Flu and Pneumonia	http://msdh.ms.gov/msdhsite/_static/14,0,199.html
World Health Organization FluNet	http://www.who.int/influenza/gisrs laboratory/flunet/en/


Appendix

















Week Ending



Sep.

0^{c.21}






Figure 7

